

University of Groningen

Bridging Binding Modes of Phosphine-Stabilized Nitrous Oxide to Zn(C₆F₅)₂

Neu, Rebecca C.; Otten, Edwin; Stephan, Douglas W.

Published in:
Angewandte Chemie International Edition

DOI:
[10.1002/ange.200905650](https://doi.org/10.1002/ange.200905650)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2009

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Neu, R. C., Otten, E., & Stephan, D. W. (2009). Bridging Binding Modes of Phosphine-Stabilized Nitrous Oxide to Zn(C₆F₅)₂. *Angewandte Chemie International Edition*, 48(51), 9709-9712.
<https://doi.org/10.1002/ange.200905650>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Supporting Information

© Wiley-VCH 2009

69451 Weinheim, Germany

**Bridging Binding Modes of Phosphine-Stabilized Nitrous Oxide to
 $\text{Zn}(\text{C}_6\text{F}_5)_2^{**}$**

*Rebecca C. Neu, Edwin Otten, and Douglas W. Stephan**

anie_200905650_sm_miscellaneous_information.pdf

General Considerations. All manipulations were carried out under an atmosphere of dry, O₂-free N₂ employing standard Schlenk-line and glovebox techniques. Solvents (pentane, hexanes, CH₂Cl₂) were dried by employing a Grubbs-type column system (Innovative Technology), degassed and stored under N₂. Cyclohexane was distilled under N₂ from Na/benzophenone while bromobenzene was distilled under N₂ from CaH₂. CD₂Cl₂ was vacuum transferred from CaH₂, degassed and stored under N₂. ^tBu₃P (Strem Chemicals), N₂O (Sigma-Aldrich; 99%) and ¹⁵N₂O (Cambridge Isotope Laboratories; 99.9%, 98.8% ¹⁵N enriched) were used as received. Zn(C₆F₅)₂ (Sigma-Aldrich; 97%) was recrystallized from toluene at -35 °C, yielding the toluene adduct, tol·Zn(C₆F₅)₂. The reagent B(C₆H₄F)₃¹ was prepared according to literature procedures. ¹H, ¹¹B, ¹³C, ¹⁵N, ¹⁹F and ³¹P NMR spectra were recorded at 25 °C, unless otherwise stated, on a Varian NMR System 400 MHz or Bruker Avance III 400 MHz spectrometer and were referenced using (residual) solvent resonances relative to SiMe₄ (¹H, ¹³C), or relative to an external standard (¹¹B: (Et₂O)BF₃, ¹⁹F: CFCI₃, ³¹P: 85% H₃PO₄, ¹⁵N: NH₃(l) via the ¹⁵N resonance of 90% formamide in DMSO-*d*₆ at 112 ppm).² Chemical shifts are reported in ppm and coupling constants as scalar values in Hz. Combustion analyses were performed in house employing a Perkin-Elmer CHN Analyzer.

^tBu₃PN₂OB(C₆H₄F)₃ (1). A 50 mL schlenk tube was charged with B(C₆H₄F)₃ (0.212 g, 0.716 mmol) and ^tBu₃P (0.145 g, 0.717 mmol) in bromobenzene (10 mL). The pale yellow solution was subjected to 3 freeze-pump-thaw cycles using liquid nitrogen to degas the solution. The solution was left stirring under an atmosphere of N₂O for 12 hours at room temperature. At this time, the solution was cloudy and pale yellow. Pentane (10 mL) was added precipitating a white solid. The solid was isolated by filtration, washed with pentane (3 x 5 mL) and dried *in vacuo* for 2 hours. Yield: 0.312 g (80 %). ¹H NMR (400 MHz, CD₂Cl₂, 25 °C): δ 7.33 (m, 6H, *o*-C₆H₄F), 6.87 (m, 6H, *m*-C₆H₄F), 1.38 (d, 27H, ³J_{H-P} = 14 Hz). ¹¹B{¹H} NMR (128 MHz, CD₂Cl₂, 25 °C): δ 6.69 (s). ¹³C{¹H} NMR (101 MHz, CD₂Cl₂, 25 °C) partial: δ 161.25 (br d, ¹J_{C-F} = 247 Hz, *p*-C₆H₄F), 135.54 (br s, *m*-C₆H₄F), 113.50 (br s, *o*-C₆H₄F), 41.31 (d, ¹J_{C-P} = 31 Hz, P{C(CH₃)₃}), 29.85 (s, P{C(CH₃)₃}). ¹⁹F NMR (376 MHz, CD₂Cl₂, 25 °C): δ -120.87 (s). ³¹P{¹H} NMR (162 MHz, CD₂Cl₂, 25 °C): δ 65.38 (s). Anal. Calcd. for C₃₀H₃₉BF₃N₂OP: C, 66.43; H, 7.25; N, 5.16. Found: C, 66.25; H, 7.27; N, 5.17 %.

^tBu₃P¹⁵N₂OB(C₆H₄F)₃ (1-¹⁵N). A J. Young NMR tube containing a solution of B(C₆H₄F)₃ (0.025 g, 0.084 mmol) and ^tBu₃P (0.017 g, 0.084 mmol) in C₆H₅Br (0.5 mL) was subjected to 3 freeze-pump-thaw cycles using liquid nitrogen to degas the solution and backfilled with ¹⁵N₂O. After standing overnight, the product was precipitated by addition of hexanes (5 mL). The product was allowed to settle and the solvent was decanted followed by washing with hexanes (2 x 1 mL). The solid was dried *in vacuo* for 1 hour. ¹H and ¹⁹F NMR spectra were identical to **1**. ¹⁵N NMR (40.6

MHz, CD₂Cl₂, 25 °C): δ 588.75 (dd, ²J_{N-P} = 19 Hz, ¹J_{N-N} = 18 Hz, PNNO), 367.61 (dd, ¹J_{N-P} = 61 Hz, ¹J_{N-N} = 18 Hz, PMNO). ³¹P{¹H} NMR (162 MHz, CD₂Cl₂, 25 °C): δ 64.27 (dd, ¹J_{P-N} = 61 Hz, ²J_{P-N} = 19 Hz).

[^tBu₃PN₂OZn(C₆F₅)₂]₂ (2). A 20 mL scintillation vial was charged with **1** (0.100 g, 0.184 mmol) and tol-Zn(C₆F₅)₂ (0.091 g, 0.185 mmol) in CH₂Cl₂ (5 mL). The solution was initially opaque but cleared after a few seconds of stirring. The reaction was left stirring for 1 hour at room temperature. At this time, the solution was cloudy. Hexanes (10 mL) was added precipitating a fine white solid. The solid was isolated by filtration, washed with hexanes (3 x 5 mL) and dried *in vacuo* for 2 hours. Crystals suitable for X-ray diffraction were grown from a layered CH₂Cl₂/pentane solution at -35 °C. Yield: 0.118 g (99 %). ¹H NMR (400 MHz, CD₂Cl₂, 25 °C): δ 1.51 (d, 27H, ³J_{H-P} = 14 Hz). ¹³C{¹H} NMR (101 MHz, CD₂Cl₂, 25 °C): δ 149.30 (br d, ¹J_{C-F} = 228 Hz, *o*-C₆F₅), 140.46 (br d, ¹J_{C-F} = 244 Hz, *p*-C₆F₅), 136.90 (br d, ¹J_{C-F} = 259 Hz, *m*-C₆F₅), 120.71 (s, *ipso*-C₆F₅), 41.65 (d, ¹J_{C-P} = 31.65, P{C(CH₃)₃}), 29.70 (s, P{C(CH₃)₃}). ¹⁹F NMR (376 MHz, CD₂Cl₂, 25 °C): δ -117.44 (m, *o*-C₆F₅), -157.71 (t, ³J_{F-F} = 19 Hz, *p*-C₆F₅), -162.64 (m, *m*-C₆F₅). ³¹P{¹H} NMR (162 MHz, CD₂Cl₂, 25 °C): δ 66.50 (s). Anal. Calcd. for C₄₈H₅₄F₂₀N₄O₂P₂Zn₂: C, 44.63; H, 4.21; N, 4.34. Found: C, 44.32; H, 4.14; N, 4.66 %.

[^tBu₃P¹⁵N₂OZn(C₆F₅)₂]₂ (2-¹⁵N). A 10 mL scintillation vial was charged with **1-¹⁵N** (0.023 g, 0.042 mmol) and tol-Zn(C₆F₅)₂ (0.021 g, 0.043 mmol) in CH₂Cl₂ (3 mL). The clear solution was left stirring for 1 hour at room temperature. At this time, hexanes (5 mL) was added precipitating a fine white solid. The solid was allowed to settle and the solvent was decanted followed by washing of the solid with hexanes (2 x 1 mL). The product was dried *in vacuo* for 1 hour. ¹H and ¹⁹F NMR spectra are identical to **2**. ¹⁵N NMR (40.6 MHz, CD₂Cl₂, 25 °C): δ 599.07 (dd, ²J_{N-P} = 9.3 Hz, ¹J_{N-N} = 18 Hz, PNNO), 317.97 (dd, ¹J_{N-P} = 54 Hz, ¹J_{N-N} = 18 Hz, PMNO). ³¹P{¹H} NMR (162 MHz, CD₂Cl₂, 25 °C): δ 65.83 (dd, ¹J_{P-N} = 54 Hz, ²J_{P-N} = 9.3 Hz).

(^tBu₃N₂OZn(C₆F₅)₂)₂Zn(C₆F₅)₂ (3). Method I. A 20 mL scintillation vial was charged with **1** (0.060 g, 0.111 mmol) and tol-Zn(C₆F₅)₂ (0.082 g, 0.167 mmol) in CH₂Cl₂ (10 mL). The clear solution was left stirring for 1 hour at room temperature. At this time, pentane (10 mL) was added precipitating a fine white solid. The product was allowed to settle and the solvent was decanted followed by washing of the solid with pentane (3 x 5 mL). The product was dried *in vacuo* for 1 hour. Yield: 0.076 g (81 %). **Method II.** A 20 mL scintillation vial was charged with **2** (0.037 g, 0.029 mmol) and tol-Zn(C₆F₅)₂ (0.014 g, 0.028 mmol) in CH₂Cl₂ (10 mL). The clear solution was left stirring for 1 hour at room temperature. At this time, pentane (10 mL) was added precipitating a fine white solid. The product was allowed to settle and the solvent was decanted followed by washing of the solid with pentane (3 x 5 mL). The product was dried *in vacuo* for 1 hour. Yield:

0.037 g (77 %). Crystals suitable for X-ray diffraction were grown from a layered CH₂Cl₂/pentane solution at -35 °C. ¹H NMR (400MHz, CD₂Cl₂, 25 °C): δ 1.46 (d, 27H, ³J_{H-P} = 14 Hz). ¹³C{¹H} NMR (101 MHz, CD₂Cl₂, 25 °C): δ 149.26 (br d, ¹J_{C-F} = 245 Hz, *o*-C₆F₅), 140.17 (br d, ¹J_{C-F} = 224 Hz, *p*-C₆F₅), 136.41 (br d, ¹J_{C-F} = 241 Hz, *m*-C₆F₅), 118.95 (s, *ipso*-C₆F₅), 41.86 (d, ¹J_{C-P} = 30 Hz, P{C(CH₃)₃}), 29.59 (s, P{C(CH₃)₃}). ¹⁹F NMR (376 MHz, CD₂Cl₂, 25 °C): δ -117.56 (m, *o*-C₆F₅), -156.73 (t, ³J_{F-F} = 19 Hz, *p*-C₆F₅), -162.42 (m, *m*-C₆F₅). ³¹P{¹H} NMR (162 MHz, CD₂Cl₂, 25 °C): δ 68.54 (s). Anal. Calcd. for C₆₀H₅₄F₃₀N₄O₂P₂Zn₃·CH₂Cl₂: C, 41.25; H, 3.18; N, 3.15. Found: C, 41.73; H, 3.39; N, 3.49 %.

(¹Bu₃PN₂OZn(C₆F₅)₂)₂Zn(C₆F₅)₂ at -75 °C. A J. Young NMR tube was charged with **3** (0.006 g, 0.004 mmol) in CD₂Cl₂ (0.5 mL). The ¹⁹F NMR spectrum was recorded at -75 °C. ¹⁹F NMR (376 MHz, CD₂Cl₂, -75 °C): δ -117.43 (br s, *o*-C₆F₅), -117.93 (br s, *o*-C₆F₅), -156.26 (t, ³J_{F-F} = 19 Hz, *p*-C₆F₅), -157.65 (t, ³J_{F-F} = 18 Hz, *p*-C₆F₅), -161.81 (br s, *m*-C₆F₅), -162.30 (br s, *m*-C₆F₅).

(¹Bu₃N₂OZn(C₆F₅)₂)₂Zn(C₆F₅)₂ (3-¹⁵N). A 10 mL scintillation vial was charged with **1-¹⁵N** (0.010 g, 0.018 mmol) and tol·Zn(C₆F₅)₂ (0.014 g, 0.028 mmol) in CH₂Cl₂ (3 mL). The clear solution was left stirring for 1 hour at room temperature. At this time, hexanes (5 mL) was added precipitating a fine white solid. The solid was allowed to settle and the solvent was decanted followed by washing with hexanes (2 x 1 mL). The solid was then dried *in vacuo* for 1 hour. ¹H and ¹⁹F NMR spectra are identical to **3**. ¹⁵N NMR (40.6 MHz, CD₂Cl₂, 25 °C): δ 595.17 (dd, ²J_{N-P} = 9.4 Hz, ¹J_{N-N} = 18 Hz, PNNO), 323.78 (dd, ¹J_{N-P} = 54 Hz, ¹J_{N-N} = 18 Hz, PMNO). ³¹P{¹H} NMR (162 MHz, CD₂Cl₂, 25 °C): δ 68.22 (dd, ¹J_{P-N} = 54 Hz, ²J_{P-N} = 9.4 Hz).

¹Bu₃PN₂O(Zn(C₆F₅)₂)₂ (4). Method I. A 20 mL scintillation vial was charged with **1** (0.064 g, 0.118 mmol) and tol·Zn(C₆F₅)₂ (0.116 g, 0.236 mmol) in CH₂Cl₂ (10 mL). The clear solution was left stirring for 1 hour at room temperature. At this time, pentane (10 mL) was added precipitating a fine white solid. The product was allowed to settle and the solvent was decanted followed by washing with pentane (3 x 5 mL). The solid was dried *in vacuo* for 1 hour. Yield: 0.099 g, (80 %). **Method II.** A 20 mL scintillation vial was charged with **2** (0.060 g, 0.046 mmol) and tol·Zn(C₆F₅)₂ (0.046 g, 0.094 mmol) in CH₂Cl₂ (8 mL). The cloudy, faintly yellow solution was left stirring for 1 hour at room temperature. At this time, hexanes (10 mL) was added precipitating a fine white powder. The solid was allowed to settle and the solvent was decanted followed by washing with hexanes (3 x 5 mL). The solid was dried *in vacuo* for 1 hour. Yield: 0.068 g (70 %). Crystals suitable for X-ray diffraction were grown from a layered CH₂Cl₂/cyclohexane solution at 25 °C. ¹H NMR (400MHz, CD₂Cl₂, 25 °C): δ 1.43 (d, 27H, ³J_{H-P} = 14 Hz). ¹³C{¹H} NMR (101 MHz, CD₂Cl₂, 25 °C): δ 149.41 (br d, ¹J_{C-F} = 224 Hz, *o*-C₆F₅), 140.97 (br d, ¹J_{C-F} = 248 Hz, *p*-C₆F₅), 137.00 (br d, 262 Hz, *m*-C₆F₅), 118.29 (s, *ipso*-C₆F₅), 42.04 (d, ¹J_{C-P} = 30 Hz, P{C(CH₃)₃}), 29.53 (s,

$\text{P}\{\text{C}(\text{CH}_3)_3\}$. ^{19}F NMR (376 MHz, CD_2Cl_2 , 25 °C): δ -117.62 (m, *o*- C_6F_5), -156.26 (t, $^3J_{\text{F-F}} = 20$ Hz, *p*- C_6F_5), -162.18 (m, *m*- C_6F_5). $^{31}\text{P}\{^1\text{H}\}$ NMR (162 MHz, CD_2Cl_2 , 25 °C): δ 71.65 (s). Anal. Calcd. for $\text{C}_{36}\text{H}_{27}\text{F}_{20}\text{N}_2\text{OPZn}_2 \cdot 0.5\text{CH}_2\text{Cl}_2$: C, 40.30; H, 2.59; N, 2.57. Found: C, 40.05; H, 2.80; N, 2.78 %.

$^t\text{Bu}_3\text{PN}_2\text{O}(\text{Zn}(\text{C}_6\text{F}_5)_2)_2$ at -75 °C. A J. Young NMR tube was charged with **4** (0.006 g, 0.006 mmol) in CD_2Cl_2 (0.5 mL). The ^{19}F NMR spectrum was recorded at -75 °C. ^{19}F NMR (376 MHz, CD_2Cl_2 , -75 °C): δ -117.61 (d, $^3J_{\text{F-F}} = 20$ Hz, *o*- C_6F_5), -118.15 (d, $^3J_{\text{F-F}} = 20$ Hz, *o*- C_6F_5), -154.85 (t, $^3J_{\text{F-F}} = 20$ Hz, *p*- C_6F_5), -155.36 (t, $^3J_{\text{F-F}} = 21$ Hz, *p*- C_6F_5), -161.10 (t, $^3J_{\text{F-F}} = 19$ Hz, 2 x *m*- C_6F_5).

$^t\text{Bu}_3\text{P}^{15}\text{N}_2\text{O}(\text{Zn}(\text{C}_6\text{F}_5)_2)_2$ (4**- ^{15}N)**. A 10 mL scintillation vial was charged with **2**- ^{15}N (0.014 g, 0.011 mmol) and $(\text{C}_6\text{F}_5)_2\text{Zn}$ -toluene (0.010 g, 0.020 mmol) in dichloromethane (3 mL). The clear solution was left stirring for 1 hour at room temperature. At this time, hexanes (5 mL) was added precipitating a fine white solid. The solid was allowed to settle and the solvent was decanted followed by washing with hexanes (2 x 1 mL). The product was dried *in vacuo* for 1 hour. ^1H and ^{19}F NMR spectra are identical to **4**. ^{15}N NMR (40.6 MHz, CD_2Cl_2 , 25 °C): δ 582.52 (dd, $^2J_{\text{N-P}} = 11$ Hz, $^1J_{\text{N-N}} = 17$ Hz, PNNO), 349.33 (dd, $^1J_{\text{N-P}} = 54$ Hz, $^1J_{\text{N-N}} = 17$ Hz, PMNO). $^{31}\text{P}\{^1\text{H}\}$ NMR (162 MHz, CD_2Cl_2 , 25 °C): δ 71.99 (dd, $^1J_{\text{P-N}} = 54$ Hz, $^2J_{\text{P-N}} = 11$ Hz).

References

1. D. Naumann, H. Butler, R. Z. Gnann, *Anorg. Allg. Chem.* **1992**, 618.
2. G. E. Martin, C. E. Hadden, *J. Nat. Prod.* **2000**, 63, 543.